

# *Xanthomonas euroxanthea* sp. nov., a new xanthomonad species including pathogenic and non-pathogenic strains of walnut

Leonor Martins<sup>1,2,\*</sup>, Camila Fernandes<sup>1,2,3</sup>, Jochen Blom<sup>4</sup>, Nay C. Dia<sup>5,6</sup>, Joël F. Pothier<sup>5</sup> and Fernando Tavares<sup>1,2,\*</sup>

## Abstract

We describe a novel species isolated from walnut (*Juglans regia*) which comprises non-pathogenic and pathogenic strains on walnut. The isolates, obtained from a single ornamental walnut tree showing disease symptoms, grew on yeast extract–dextrose–carbonate agar as mucoid yellow colonies characteristic of *Xanthomonas* species. Pathogenicity assays showed that while strain CPBF 424<sup>T</sup> causes disease in walnut, strain CPBF 367 was non-pathogenic on walnut leaves. Biolog GEN III metabolic profiles disclosed some differences between strains CPBF 367 and CPBF 424<sup>T</sup> and other xanthomonads. Multilocus sequence analysis with seven housekeeping genes (*fyuA*, *gyrB*, *rpoD*, *atpD*, *dnaK*, *efp*, *glnA*) grouped these strains in a distinct cluster from *Xanthomonas arboricola* pv. *juglandis* and closer to *Xanthomonas prunicola* and *Xanthomonas arboricola* pv. *populi*. Average nucleotide identity (ANI) analysis results displayed similarity values below 93% to *X. arboricola* strains. Meanwhile ANI and digital DNA–DNA hybridization similarity values were below 89 and 50% to non-*arboricola* *Xanthomonas* strains, respectively, revealing that they do not belong to any previously described *Xanthomonas* species. Furthermore, the two strains show over 98% similarity to each other. Genomic analysis shows that strain CPBF 424<sup>T</sup> harbours a complete type III secretion system and several type III effector proteins, in contrast with strain CPBF 367, shown to be non-pathogenic in plant bioassays. Taking these data altogether, we propose that strains CPBF 367 and CPBF 424<sup>T</sup> belong to a new species herein named *Xanthomonas euroxanthea* sp. nov., with CPBF 424<sup>T</sup> (=LMG 31037<sup>T</sup>=CCOS 1891<sup>T</sup>=NCPPB 4675<sup>T</sup>) as the type strain.

## INTRODUCTION

The taxonomy of the genus *Xanthomonas* has been extensively studied and categorized over last century, raising some disputes regarding classification and nomenclature of its members, largely due to the wide plant host diversity associated with the genus. Recorded *Xanthomonas* hosts include numerous economically relevant crop species, severely impacting agricultural productivity worldwide [1].

Introduction of molecular methods such as DNA–DNA hybridization allowed the first big reclassification within the genus *Xanthomonas* [2]. Recently, studies relying on multi-locus sequence analysis (MLSA), whole-genome average

nucleotide identity (ANI) and biochemical analysis have proved fruitful in the reclassification of *Xanthomonas* species and description amendments of several strains [3, 4]. These techniques have allowed us to reach a deeper discriminatory insight into the genetic diversity of *Xanthomonas* at the intrasubspecific level, consequently leading to the proposal of novel species [5–7].

Increasing reports of non-pathogenic *Xanthomonas* strains, such as *Xanthomonas arboricola* isolates recovered from asymptomatic and symptomatic host tissues, points to the uncertainty of several *Xanthomonas* strains regarding their host-associated lifestyle and likely host diversity [8–10].

**Author affiliations:** <sup>1</sup>CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO, Laboratório Associado, Universidade do Porto, Vairão, Portugal; <sup>2</sup>FCUP, Faculdade de Ciências, Departamento de Biologia, Universidade do Porto, Rua do Campo Alegre s/n, Porto, Portugal; <sup>3</sup>INIAV, Instituto Nacional de Investigação Agrária e Veterinária, Av. da República, Quinta do Marquês, Oeiras, Portugal; <sup>4</sup>Bioinformatics and Systems Biology, Justus-Liebig-University Giessen, Giessen, Germany; <sup>5</sup>Environmental Genomics and Systems Biology, Institute of Natural Resource Sciences, Zurich University of Applied Sciences (ZHAW), Wädenswil, Switzerland; <sup>6</sup>Molecular Plant Breeding, Institute of Agricultural Sciences, ETH Zurich, Zurich, Switzerland.

\*Correspondence: Leonor Martins, leonor.martins@cibio.up.pt; Fernando Tavares, ftavares@fc.up.pt

**Keywords:** new species; pathogenicity; walnut; *Xanthomonas*.

**Abbreviations:** ANI, average nucleotide identity; BLASTn, nucleotide Basic Local Alignment Search Tool; dDDH, digital DNA–DNA hybridization; MLSA, multilocus sequence analysis; T3E, type III effectors; T3SS, type III secretion system; WBB, walnut bacterial blight; YDC, yeast extract–dextrose–carbonate.

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain CPBF 424<sup>T</sup> is MT036365. The GenBank accession numbers for genome sequences of CPBF 367 and CPBF 424<sup>T</sup> are UNRN00000000.1 and UIHB00000000.1, respectively.

One supplementary figure and two supplementary tables are available with the online version of this article.

004386 © 2020 The Authors



This is an open-access article distributed under the terms of the Creative Commons Attribution NonCommercial License.

**Table 1.** Strains used for pathogenicity tests and Biolog assays

Strain	Species	Host of isolation	Geographic origin	Year of isolation
CPBF 367*	<i>X. euroxanthea</i>	<i>Juglans regia</i>	Loures, Portugal	2016
CPBF 424 <sup>T</sup> *	<i>X. euroxanthea</i>	<i>Juglans regia</i>	Loures, Portugal	2016
CPBF 427*	<i>X. arboricola</i> pv. <i>juglandis</i>	<i>Juglans regia</i>	Loures, Portugal	2016
CPBF 1521*	<i>X. arboricola</i> pv. <i>juglandis</i>	<i>Juglans regia</i>	Loures, Portugal	2014
CPBF 1480	<i>X. arboricola</i> pv. <i>juglandis</i>	<i>Juglans regia</i>	Azeitão, Portugal	2014
LMG 747 <sup>T</sup>	<i>X. arboricola</i> pv. <i>juglandis</i>	<i>Juglans regia</i>	New Zealand	1956
CFBP 3123 <sup>PT</sup>	<i>X. arboricola</i> pv. <i>populi</i>	<i>Populus</i> × <i>euroamericana</i> cv. <i>robusta</i>	The Netherlands	1979

\*Other known names for these strains: CPBF 367=LMG 31036=CCOS 1890; CPBF 424<sup>T</sup>=LMG 31037<sup>T</sup>=CCOS 1891<sup>T</sup>=NCPBP 4675<sup>T</sup>; CPBF 427=LMG 31039=CCOS 1893; CPBF 1521=LMG 31040=CCOS 1894=NCPBP 4676. (CPBF, Portuguese Collection of Phytopathogenic Bacteria in Oeiras, Portugal; BCCM/LMG, the Belgian Coordinated Collections of Microorganisms/ LMG Bacteria Collection in Gent, Belgium; CCOS, Culture Collection of Switzerland in Wädenswil, Switzerland; NCPBP, National Collection of Plant Pathogenic Bacteria in York, UK; CFBP, French Collection of Plant associated Bacteria, France).

Interestingly, *X. arboricola* is the species with most non-pathogenic strains within the genus *Xanthomonas* [11]. Characterization of several of these non-pathogenic *X. arboricola* strains has revealed that they are phylogenetically distant from the disease-causing strains that were isolated from their specific hosts [8]. Furthermore, plants are host to a consortium of both distant and genetically similar bacteria, displaying different host-associated phenotypes. It has been suggested that the sympatry of pathogenic and non-pathogenic strains may favour horizontal gene transfer events, likely leading to the emergence of new pathogenic lineages [8, 11].

The advent of increasingly powerful genomic tools and improved bacterial genome assemblers and accurate annotation has led to a more confident examination of bacterial evolution and phylogeny, allowing microbiologists and plant pathologists to assign specific differences to the existing taxonomy, even at the pathovar level. A pathovar of a pathogenic species refers to an infrasubspecific group that includes strains with the same host range [12, 13]. To the best of our knowledge, all *Xanthomonas arboricola* pv. *juglandis* strains described previously have been isolated from diseased walnut trees.


Furthermore, MLSA results revealed that several *X. arboricola* strains isolated from walnut, including both pathogenic and non-pathogenic strains, do not cluster with the members of *juglandis* pathovar [9, 14]. Indeed, a previous study has reported the identification of two *X. arboricola* strains isolated from symptomatic leaves of a pecan tree (*Carya illinoensis*), which are pathogenic to both pecan and walnut trees [14]. These data raise the need to understand how the genetic makeup of plant bacterial pathogens, such as *Xanthomonas*, links with a diversifying range of plant hosts. Walnut (*Juglans* species) is an economically important tree crop worldwide, with the Persian walnut (*Juglans regia* L.) being the main cultivated species for walnut fruits and timber production [15–17]. *X. arboricola* pv. *juglandis* is a major threat to walnut orchards and acknowledged as the etiological agent of walnut

bacterial blight (WBB) and other walnut diseases such as brown apical necrosis [18] and vertical oozing canker [19], altogether causing high yield losses [16].

In this study we characterize two novel strains, CPBF 367 and CPBF 424<sup>T</sup>, isolated from asymptomatic buds of one walnut tree showing WBB symptoms. Genotyping and comparative genomics indicate that these strains, showing a distinct pathogenicity phenotype in walnut, belong to the genus *Xanthomonas*, although they could not be assigned to any of the known species of this genus. This work gathers phenotypic, genotypic and genomic evidence to support that strains CPBF 367 and CPBF 424<sup>T</sup> are members of a new species of the genus *Xanthomonas*.

## ISOLATION AND GROWTH OF BACTERIA

Several isolates were obtained over three consecutive years (2014–2016) from an isolated ornamental walnut tree (*Juglans regia*) showing bacterial blight symptoms in leaves and fruits, located in a municipal garden in the city of Loures, Portugal (Fig. S1, available in the online version of this article). Distinct organs were sampled including symptomatic leaves (CPBF 1521 in 2014) and asymptomatic buds (CPBF 367, CPBF 424<sup>T</sup>, CPBF 426 and CPBF 427 in 2016). The procedure for preparation of plant material for bacteria isolation has been described previously [20]. Briefly, the excised plant material was disinfected by immersion in 70% ethanol followed by washing with sterile distilled water (SDW) and then macerated in extraction bags with 5 ml SDW [21]. Suspension and correspondent dilutions were streaked on yeast extract–dextrose–carbonate (YDC) agar [22] and incubated at 26±2 °C for 4 days. Single characteristic mucoid yellow colonies of *Xanthomonas* were streaked on fresh nutrient agar (NA) medium to ensure purity. Selected isolates (CPBF 424<sup>T</sup> and CPBF 367) were stored at –80 °C in cryovials containing LB medium (Difco) and glycerol to a final concentration of 30%. The strains were deposited in international bacterial collections with the accession



**Fig. 1.** Walnut plantlets leaves 7 days post-inoculation with (a) isolate CPBF 424<sup>T</sup> showing walnut bacteria blight (WBB) symptoms, (b) isolate CPBF 367 no WBB symptoms were observed, (c) *Xanthomonas arboricola* pv. *juglandis* LMG 747<sup>T</sup> and (d) *X. arboricola* pv. *juglandis* strain CPBF 1480, used as positive controls, show characteristic WBB symptoms.

numbers CPBF 367 (LMG 31036=CCOS 1890) and CPBF 424<sup>T</sup> (=LMG 31037<sup>T</sup>=CCOS 1891<sup>T</sup>=NCPBPB 4675<sup>T</sup>). Isolates CPBF 367 and CPBF 424<sup>T</sup> were grown in YDC medium for analysis of colony morphology and pigment production for a 10 day period. The bacterial strains used throughout this study are disclosed in Table 1.

## PATHOGENICITY ASSAYS

Plantlets used in the pathogenicity tests were grown from seeds collected from *Juglans regia* cv. Hartley on the same day. After 30 days of cold stratification treatment at 3–5 °C to break dormancy, seeds were sown in sterilized sand substrate and germinated over 60 days at alternated temperatures, 16 h day at 30 °C and 8 h night at 20 °C [23]. Walnut plantlets were then maintained in a climatic chamber under controlled environmental conditions of 16 h photoperiod (16 h of light at 24 °C and 8 h of darkness at 18 °C). Pathogenicity determination of strains CPBF 424<sup>T</sup> and CPBF 367 was carried out as previously described [20]. Briefly, inoculum suspensions of approximately 10<sup>8</sup> c.f.u. ml<sup>-1</sup> were prepared with SDW. Plantlets with at least four young leaves expanded were inoculated by spraying with a manual atomizer until runoff. SDW was used for negative control, while *X. arboricola* pv. *juglandis* strains LMG 747<sup>T</sup> and CPBF 1480 were used as positive controls of infection. Three replicates were included for each strain tested. Symptoms were registered after 7 days and further followed for over 5 weeks. Necrotic bacterial spots were observed 7 days post inoculation on leaves of plants inoculated with walnut isolate CPBF 424<sup>T</sup>, and with positive-control strains LMG 747<sup>T</sup> and CPBF 1480 (Fig. 1). Walnut isolate CPBF 367 and negative controls did not produce any disease symptoms on walnut plantlets. In order to fulfil Koch's postulates, typical *Xanthomonas* yellow mucoid bacterial colonies were re-isolated from the symptoms and confirmed by sequencing analysis of *gyrB* and *fyuA*.

## PHENOTYPIC CHARACTERIZATION

Phenotypic characterization was carried out with Biolog GEN III MicroPlates (Biolog) according to the manufacturer's instructions, on walnut isolates CPBF 367, CPBF 424<sup>T</sup>, CPBF 427, CPBF 1521 and reference strains *X. arboricola* pv. *juglandis* LMG 747<sup>T</sup> and *X. arboricola* pv. *populi* CFBP 3123<sup>PT</sup>. For all strains except CFBP 3123<sup>PT</sup>, three MicroPlates were assayed on different dates. Selected strains were first grown on solid NYGA medium (5 g l<sup>-1</sup> peptone, 3 g l<sup>-1</sup> yeast extract, 20 g l<sup>-1</sup> glycerol and 15 g l<sup>-1</sup> agar) for 48 h at 28 °C. The strains were subsequently grown on solid Biolog Dehydrated Growth agar for 24 h at 28 °C. Fresh colonies were then transferred into Inoculating Fluid A vials using a cotton-tipped swab. The density of the inoculum was checked and adjusted to a transmittance of 95–98% using a turbidimeter. A total of 100 µl prepared inoculum was then dispensed into each well of the Biolog MicroPlate. MicroPlates were then incubated at 28 °C and read using a MicroStation 2 Reader (Biolog) at 24, 48, 72, 120, 192 and 240 h.

Considering compound metabolism, values 160% superior to the negative control (water) value were considered positive, while those below 130% the negative control were taken as negative. Values in between were considered borderline [10]. The data measured at 72 h was shown to provide the most consistent results between replicates and thus was selected for further analysis. Isolates CPBF 367 and CPBF 424<sup>T</sup> seemed to metabolize mucic acid and D-saccharic acid, in contrast to *X. arboricola* pv. *juglandis* CPBF 427, CPBF 1521 and LMG 747<sup>T</sup>. Additionally, comparisons with results from other *Xanthomonas* strains [4–6] revealed differences in metabolizing D-salicin, sucrose, acetic acid, melibiose, formic acid, D-arabitol, D-glucose-6-phosphate, maltose and gentiobiose (Table 2). The metabolization of some substrates was inconclusive due to the inconsistencies observed between replicates.


**Table 2.** Biolog profiles of *Xanthomonas euroxantha* strains in comparison with members of *Xanthomonas arboricola* pv. *juglandis* and other related *Xanthomonas* species

+, Positive; -, negative; BL, border line; (+), weak positive; NA, not analysed.

Test	<i>X. euroxantha</i> CPBF 367	<i>X. euroxantha</i> CPBF 424 <sup>†</sup>	<i>X. arboricola</i> pv. <i>juglandis</i> CPBF 427	<i>X. arboricola</i> pv. <i>juglandis</i> CPBF 1521	<i>X. arboricola</i> pv. <i>juglandis</i> LMG 747 <sup>†</sup>	<i>X. arboricola</i> pv. <i>populi</i> CFBP 3123 <sup>††</sup>	<i>X. floridensis</i> <sup>*</sup>	<i>X. nasturtii</i> <sup>*</sup>	<i>X. campestris</i> pv. <i>campestris</i> <sup>*</sup>	<i>X. axonopodis</i> pv. <i>phaseoli</i> <sup>*</sup>	<i>X. prunicola</i> <sup>†</sup>	<i>X. arboricola</i> <sup>‡</sup>
D-Salicin	-	-	-	-	-	+	+	-	-	-	N/A	N/A
Mucic acid	+	+	-	-	-	+	+	+	+	+	N/A	N/A
D-Saccharic acid	+	+	-	-	-	+	+	+	+	+	N/A	-
Vancomycin	+/BL	+/BL	-	-	-/BL	-	-	-	-	-	N/A	N/A
Sucrose	+	+	+	+	-	+	+	+	+	+	+	N/A
Acetic acid	+	+	+	+	+	+	+	+	+	+	-	+
Melibiose	+	+	+	+	+	+	+	(+)/+	+	+	-	N/A
Formic acid	+	+	+	+	+	+	(+)	-	-	-	-	-
D-Arabitol	-	-	-	-	-	BL	+	+	-	(+)	N/A	N/A
D-Glucose-6-phosphate	-	-	-	-	-	+	-	-	-	-	-	N/A
Maltose	+	+	+	+	+	+	+	+	-	-	+	+
Gentiobiose	+	+	+	+	+	+	+	(+)	(+)	-	+	+

\*Vicente et al. 2017 [5] (*X. floridensis* WHRI 8844, WHRI 8846A, WHRI 8848<sup>†</sup>, WHRI 8851; *X. nasturtii* WHRI 8853<sup>†</sup>, WHRI 8930A; *X. campestris* pv. *campestris*: WHRI 1279A); *X. axonopodis* pv. *phaseoli* WHRI 1925C).†López et al. 2018 [6] (*X. prunicola* CFBP 8353<sup>†</sup>, CFBP 8354, CFBP 8355).‡Vauterin et al. 1995 [2] (*X. arboricola* pv. *corylina* LMG 688; LMG 689<sup>†</sup>, LMG 8658; LMG 8660; *X. arboricola* pv. *juglandis* LMG 747<sup>†</sup>, LMG 8047; *X. arboricola* pv. *poinsetticola* LMG 5403; *X. arboricola* pv. *populi* LMG 12141<sup>††</sup>; *X. arboricola* pv. *pruni* LMG 852<sup>†</sup>, LMG 8680).





**Fig. 3.** Average nucleotide identity (ANI) between novel isolates CPBF 367 and CPBF 424<sup>T</sup> (in bold) and 20 different *Xanthomonas* genomes using EDGAR. Pairwise ANI values are shown in percentages and indicated within the cells of the heat-map. Accession numbers of sequences are available in Table S1.


367 and CPBF 426 was carried out as described for CPBF 424<sup>T</sup> [25]. Genomes of strains CPBF 367 and CPBF 424<sup>T</sup> were approximately 4.96 and 4.90 Mbp, respectively, and comprised 22 and 10 contigs, respectively. High-quality draft genomes of strains CPBF 367 and CPBF 424<sup>T</sup> were deposited in GenBank under accession numbers UNRN00000000.1 and UIHB00000000.1, respectively.

ANI and digital DNA–DNA hybridization (dDDH) values between CPBF 367 and CPBF 424<sup>T</sup>, and genomes of *X. arboricola* and *Xanthomonas* species were calculated using the EDGAR version 2.0 platform [26, 27] and formula 2 of the Genome-to-Genome Distance Calculator (GGDC 2.1; <http://ggdc.dsmz.de/ggdc.php>) [28]. ANI analysis included the genomes of 10 *X. arboricola* strains, belonging to eight distinct pathovars, and 10 other *Xanthomonas* species (Table S1). ANI values of strains CPBF 367 and CPBF 424<sup>T</sup> were less than 93% with *X. arboricola* species and less than 89% with non-arboricola *Xanthomonas* species (Fig. 3). These values reside below the threshold of 95–96% commonly considered for representatives of the same species [29]. Furthermore, these CPBF 367 and CPBF 424<sup>T</sup> strains share 98% ANI with each other (Fig. 3). The dDDH analysis considered 20 strains representing different species of the genus *Xanthomonas* and revealed similarities values below the 70% threshold (Table S2) considered for delimitation of new species [28]. Taken together, the results of ANI and dDDH analyses suggest that these strains belong to a new *Xanthomonas* species.

## TYPE III SECRETION SYSTEM (T3SS) AND TYPE III EFFECTORS (T3E)

The prediction of type III secretion system (T3SS) and type III effectors (T3E) homologs was carried out with BLASTn analysis against previously characterized T3SS and T3Es of the *Xanthomonas* group [30–32]. The criteria applied as threshold was a minimal *e*-value of  $1e^{-10}$ , query length similarity cut-off

≥40% and sequence identity with a cut-off ≥70% (Fig. 4). The walnut pathogenic strain CPBF 424<sup>T</sup> has a T3SS gene profile similar to the strains *X. arboricola* pv. *juglandis* CPBF 427 and CPBF 1521 and possesses gene homologs described for pathogenic strains of *X. arboricola* [10, 33–35]. In contrast, strains CPBF 367 and CPBF 426 lack the majority of the T3SS- and T3E-encoding genes (Fig. 4), resembling the T3SS and T3E profile of atypical non-pathogenic strains of *Xanthomonas* species described previously [33, 35]. The absence of some of the components from the Hrp2 family and of genes that are part



**Fig. 4.** Distribution of protein homologs of the type III secretion system (T3SS) and type III effectors (T3Es) among the *Xanthomonas euroxantha* strains (CPBF 367, CPBF 424<sup>T</sup>, CPBF 426) and *Xanthomonas arboricola* pv. *juglandis* (CPBF 427 and CPBF 1521) strains based on BLASTn analysis. Dark blue corresponds to the presence of T3SS protein homologous, and dark green corresponds to the presence of T3E homologous. The light blue and green indicate absence of T3SS and T3E homologous, respectively. The threshold criteria applied were minimal *e*-value of  $1e^{-10}$ , query length similarity cut-off ≥40% and sequence identity cut-off ≥70%.

of the macromolecular structure of the T3SS may explain the non-pathogenic phenotype of CPBF 367, even while harbouring regulators genes of T3SS, such as *hrpX* and *hrpG* [36, 37]. Additionally, the fact that strain CPBF 426 is deficient for most T3SS genes, suggests that this strain may not be pathogenic as shown for CPBF 367. Intriguingly, CPBF 424<sup>T</sup> seems to have a functional T3SS and holds a narrow gene repertoire of T3E homologs in comparison with other *X. arboricola* pv. *juglandis* strains [9, 30, 33]. This differs markedly from *X. arboricola* pv. *juglandis* CPBF 427 and CPBF 1521 and strains CPBF 367 and CPBF 426. Although the pathogenic strain CPBF 424<sup>T</sup> possesses homologs for eight known effectors, from which only three were identified in non-pathogenic *X. euroxanthea* (CPBF 367 and CPBF 426), no homologs were found for most of the genes assigned as important for pathogenicity of *Xanthomonas* species, namely *avrBs2*, *avrBs2*, *avrBs3*, *avrXccA1*, *avrXccA2*, *xopAW*, *xopC*, *xopG*, *xopK*, *xopL*, *xopN*, *xopQ*, *xopV*, *xopX*, *xopZ* and *xopZ1*. All the aforementioned genes, with the exception of *xopZ2*, were present in both *X. arboricola* pv. *juglandis* CPBF 427 and CPBF 1521 strains (Fig. 4). The genomic features of this new species and the pathogen phenotype of strain CPBF 424<sup>T</sup> in walnut make these strains particularly appealing to elucidate the evolutionary hypothesis of pathogenicity in walnut, to uncover the genetic footprints of adaptation and to address speciation in *Xanthomonas*.

## DESCRIPTION OF XANTHOMONAS EUROXANTHEA SP. NOV.

*Xanthomonas euroxanthea* (eu.ro.xan.the'a. N.L. fem. adj. *euroxanthea* referring to the EuroXanth COST Action CA 16107, the EU-funded network in which the isolates of this species were characterized).

Cells are Gram-stain-negative straight rods and form colonies that are yellow, circular, smooth, mucoid and slightly convex with entire margins when grown on YDC agar for 2 days. Regarding carbon source metabolism, *X. euroxanthea* is positive for 34 substrates (dextrin, maltose, trehalose, cellobiose, gentiobiose, sucrose, melibiose, *N*-acetyl-D-glucosamine,  $\alpha$ -D-glucose, D-mannose, D-fructose, D-galactose, L-fucose, glycerol, gelatin, glycyl-L-proline, L-alanine, L-glutamic acid, L-serine, mucic acid, D-saccharic acid, methyl pyruvate, L-lactic acid, citric acid,  $\alpha$ -keto-glutaric acid, L-malic acid, bromo-succinic acid, Tween 40,  $\alpha$ -hydroxy-butyric acid,  $\alpha$ -keto-butyric acid, acetoacetic acid, propionic acid, acetic acid, formic acid) and negative for 25 compounds (turanose, stachyose, raffinose, methyl  $\beta$ -D-glucoside, D-salicin, *N*-acetyl- $\beta$ -D-mannosamine, *N*-acetyl-D-galactosamine, *N*-acetyl-neuraminic acid, 3-methyl glucose, L-rhamnose, D-sorbitol, D-arabitol, myo-inositol, D-glucose-6-phosphate, D-aspartic acid, D-serine, L-arginine, L-pyroglutamic acid, D-gluconic acid, quinic acid, *p*-hydroxy-phenylacetic acid, D-lactic acid methyl ester, D-malic acid,  $\gamma$ -amino-butyric acid and  $\beta$ -hydroxy-D,L-butyric acid).

This species includes non-pathogenic (CPBF 367) and pathogenic (CPBF 424<sup>T</sup>) strains. The type strain is CPBF 424<sup>T</sup> (LMG 31037<sup>T</sup>=CCOS 1891<sup>T</sup>=NCCPB 4675<sup>T</sup>), isolated from an

asymptomatic bud of an isolated ornamental diseased walnut tree (*J. regia*) tree growing in a municipal garden of Loures, Portugal. The high-quality draft genome of the type strain is characterized by a size of 4.8 Mbp and a G+C content of 65.9 mol%. The GenBank accession number of the 16S rRNA gene sequence of strain CPBF 424<sup>T</sup> is MT036365 and its draft genome sequence accession number is GCA\_900476395.

### Funding information

This research was co-financed by the European Structural and Investment Funds (ESIFs) through COMPETE 2020, and by National Funds through FCT- Fundação para a Ciência e Tecnologia, within the framework of the project EVOXANT (PTDC/BIA-EVF/3635/2014-POCI-01-0145-FEDER-016600). C.F. and L.M. were supported by fellowships from FCT (SFRH/BD/95913/2013; SFRH/BD/137079/2018, respectively). N.C.D. was supported by grant no IZCOZO\_177064 from the Swiss National Foundation (SNF) for Scientific Research. The EDGAR platform is financially supported by the BMBF grant FKZ031A533 within the de.NBI network. This article is based upon work from COST Action CA16107 EuroXanth, supported by COST (European Cooperation in Science and Technology).

### Acknowledgements

We thank Professor Dr Jean-Louis Charlet, Professor Dr Bernhard Schink and Professor Aharon Oren for advice regarding the nomenclature, and Christian Kunkel from the Culture Collection of Switzerland (CCOS AG, Wädenswil, Switzerland) for his support during the phenotypic characterization.

### Conflicts of interest

The authors declare that there are no conflicts of interest.

### References

- Vauterin L, Rademaker J, Swings J. Synopsis on the taxonomy of the genus *Xanthomonas*. *Phytopathology* 2000;90:677–682.
- Vauterin L, Hoste B, Kersters K, Swings J. Reclassification of *Xanthomonas*. *Int J Syst Bacteriol* 1995;45:472–489.
- da Gama MAS, Barbosa MAG, de Farias ARG, da Silva Júnior WJ, Mariano R de LR. Taxonomic repositioning of *Xanthomonas campestris* pv. *viticola* (Nayudu 1972) Dye 1978 as *Xanthomonas citri* pv. *viticola* (Nayudu 1972) Dye 1978 comb. nov. and emendation of the description of *Xanthomonas citri* pv. *anacardii* to include pigmented isolates pathogenic to cashew plant. *Phytopathology* 2018;108:1143–1153.
- Constantin EC, Cleenwerck I, Maes M, Baeyen S, Van Malderghem C et al. Genetic characterization of strains named as *Xanthomonas axonopodis* pv. *dieffenbachiae* leads to a taxonomic revision of the *X. axonopodis* species complex. *Plant Pathol* 2016;65:792–806.
- Vicente JG, Rothwell S, Holub EB, Studholme DJ. Pathogenic, phenotypic and molecular characterisation of *Xanthomonas nasturtii* sp. nov. and *Xanthomonas floridensis* sp. nov., new species of *Xanthomonas* associated with watercress production in Florida. *Int J Syst Evol Microbiol* 2017;67:3645–3654.
- López MM, Lopez-Soriano P, Garita-Cambronero J, Beltrán C, Taghouti G et al. *Xanthomonas prunicola* sp. nov., a novel pathogen that affects nectarine (*Prunus persica* var. *nectarina*) trees. *Int J Syst Evol Microbiol* 2018;68:1857–1866.
- Triplett LR, Verdier V, Campillo T, Van Malderghem C, Cleenwerck I et al. Characterization of a novel clade of *Xanthomonas* isolated from rice leaves in Mali and proposal of *Xanthomonas maliensis* sp. nov. *Antonie van Leeuwenhoek* 2015;107:869–881.
- Jacques M-A, Arlat M, Boulanger A, Boureau T, Carrère S et al. Using Ecology, Physiology, and Genomics to Understand Host Specificity in *Xanthomonas*. *Annu Rev Phytopathol* 2016;54:163–187.
- Essakhi S, Cesbron S, Fischer-Le Saux M, Bonneau S, Jacques M-A et al. Phylogenetic and variable-number tandem-repeat analyses identify nonpathogenic *Xanthomonas arboricola* lineages lacking

- the canonical type III secretion system. *Appl Environ Microbiol* 2015;81:5395–5410.
10. Garita-Cambronero J, Palacio-Bielsa A, López MM, Cubero J. Comparative genomic and phenotypic characterization of pathogenic and non-pathogenic strains of *Xanthomonas arboricola* reveals insights into the infection process of bacterial spot disease of stone fruits. *PLoS One* 2016;11:e0161977.
  11. Merda D, Bonneau S, Guimbaud J-F, Durand K, Brin C et al. Recombination-prone bacterial strains form a reservoir from which epidemic clones emerge in agroecosystems. *Environ Microbiol Rep* 2016;8:572–581.
  12. Yeates GW. A proposed nomenclature and classification for plant pathogenic bacteria. *New Zeal J Agric Res* 1978;21:153–177.
  13. Schulze-Lefert P, Panstruga R. A molecular evolutionary concept connecting nonhost resistance, pathogen host range, and pathogen speciation. *Trends Plant Sci* 2011;16:117–125.
  14. Fernandes C, Sousa R, Tavares F, Cruz L. First Report of *Xanthomonas arboricola* causing bacterial blight on pecan trees in Portugal. *Plant Disease* 2018;102:2632.
  15. Lamichhane JR. *Xanthomonas arboricola* diseases of stone fruit, almond, and walnut trees: Progress toward understanding and management. *Plant Dis* 2014;98:1600–1610.
  16. Leslie C, Uratsu S, McGranahan G, Dandekar A. Agrobacterium protocols volume 2. Springer 2006.
  17. Belisario A, Santori A, Potente G, Fiorin A, Saphy B et al. Brown apical necrosis (BAN): a fungal disease causing fruit drop of english walnut. In: Neil MC (editor). *Vi International Walnut Symposium (Acta Horticulturae. Leuven 1: International Society of Horticultural Science; 2010. pp. 449–452.*
  18. Moragrega C, Matias J, Aletà N, Montesinos E, Rovira M. Apical necrosis and premature drop of Persian (English) walnut fruit caused by *Xanthomonas arboricola* pv. *juglandis*. *Plant Dis* 2011;95:1565–1570.
  19. Hajri A, Meyer D, Delort F, Guillaumès J, Brin C et al. Identification of a genetic lineage within *Xanthomonas arboricola* pv. *juglandis* as the causal agent of vertical oozing canker of Persian (English) walnut in France. *Plant Pathol* 2010;59:1014–1022.
  20. Fernandes C, Albuquerque P, Cruz L, Tavares F. Genotyping and epidemiological metadata provides new insights into population structure of *Xanthomonas* isolated from walnut trees.
  21. Fernandes C, Albuquerque P, Sousa R, Cruz L, Tavares F. Multiple DNA markers for identification of *Xanthomonas arboricola* pv. *juglandis* isolates and its direct detection in plant samples. *Plant Dis* 2017;101:858–865.
  22. Stolp H, Starr MP. Bacteriophage reactions and speciation of Phytopathogenic *Xanthomonads*. *J Phytopathol* 1964;51:442–478.
  23. International Seed Testing Association. International rules for seed testing rules. *Seed Sci Technol* 1999;27.
  24. Fischer-Le Saux M, Bonneau S, Essakhi S, Manceau C, Jacques M-A. Aggressive emerging pathovars of *Xanthomonas arboricola* represent widespread epidemic clones distinct from poorly pathogenic strains, as revealed by multilocus sequence typing. *Appl Environ Microbiol* 2015;81:4651–4668.
  25. Fernandes C, Blom J, Pothier JF, Tavares F. High-quality draft genome sequence of *Xanthomonas* sp. strain CPBF 424, a Walnut-Pathogenic strain with atypical features. *Microbiol Resour Announc* 2018;7.
  26. Blom J, Albaum SP, Doppmeier D, Pühler A, Vorhölter F-J et al. EDGAR: a software framework for the comparative analysis of prokaryotic genomes. *BMC Bioinformatics* 2009;10:154.
  27. Blom J, Kreis J, Spänig S, Juhre T, Bertelli C et al. EDGAR 2.0: an enhanced software platform for comparative gene content analyses. *Nucleic Acids Res* 2016;44:W22–W28.
  28. Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 2013;14:60.
  29. Richter M, Rosselló-Móra R. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* 2009;106:19126–19131.
  30. Hajri A, Pothier JF, Fischer-Le Saux M, Bonneau S, Poussier S et al. Type three effector gene distribution and sequence analysis provide new insights into the pathogenicity of plant-pathogenic *Xanthomonas arboricola*. *Appl Environ Microbiol* 2012;78:371–384.
  31. Ryan RP, Vorhölter F-J, Potnis N, Jones JB, Van Sluys M-A et al. Pathogenomics of *Xanthomonas*: understanding bacterium-plant interactions. *Nat Rev Microbiol* 2011;9:344–355.
  32. White FF, Potnis N, Jones JB, Koebnik R. The type III effectors of *Xanthomonas*. *Mol Plant Pathol* 2009;10:749–766.
  33. Cesbron S, Briand M, Essakhi S, Gironde S, Boureau T et al. Comparative genomics of pathogenic and nonpathogenic strains of *Xanthomonas arboricola* unveils molecular and evolutionary events linked to pathoadaptation. *Front Plant Sci* 2015;6:1126.
  34. Garita-Cambronero J, Palacio-Bielsa A, Cubero J. *Xanthomonas arboricola* pv. *pruni*, causal agent of bacterial spot of stone fruits and almond: its genomic and phenotypic characteristics in the *X. arboricola* species context. *Mol Plant Pathol* 2018;19:2053–2065.
  35. Garita-Cambronero J, Palacio-Bielsa A, López MM, Cubero J. Pan-genomic analysis permits differentiation of virulent and non-virulent strains of *Xanthomonas arboricola* that cohabit *Prunus* spp. and elucidate bacterial virulence factors. *Front Microbiol* 2017;8:1–17.
  36. Guo Y, Figueiredo F, Jones J, Wang N. HrpG and HrpX play global roles in coordinating different virulence traits of *Xanthomonas axonopodis* pv. *citri*. *Mol Plant Microbe Interact* 2011;24:649–661.
  37. Jacobs JM, Pesce C, Lefeuvre P, Koebnik R. Comparative genomics of a cannabis pathogen reveals insight into the evolution of pathogenicity in *Xanthomonas*. *Front Plant Sci* 2015;6:431.

### Five reasons to publish your next article with a Microbiology Society journal

1. The Microbiology Society is a not-for-profit organization.
2. We offer fast and rigorous peer review – average time to first decision is 4–6 weeks.
3. Our journals have a global readership with subscriptions held in research institutions around the world.
4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at [microbiologyresearch.org](http://microbiologyresearch.org).